

# Monitoring for Insertional Oncogenesis and Clonality of Hematopoiesis after Gene Therapy

## Investigators

Fabio Candotti M.D.	NHGRI
Donald Kohn, M.D.	CHLA
Greg Podsakoff, M.D.	CHLA
Christof von Kalle, M.D.	U of Cincinnati
Kenneth Weinberg, M.D.	CHLA
Linda Forsyth, M.D.	FDA
Carolyn Wilson, Ph.D.	FDA
IRB members and staff	CHLA

# **Monitoring for Insertional Oncogenesis and Clonality of Hematopoiesis after Gene Therapy**

- **Theoretical risk of insertional oncogenesis due to:**

**RCR**

**Random or pseudo-random insertional mutagenesis**

- **Insertional mutagenesis could activate proto-oncogenes, inactivate tumor suppressor genes, or activate genes that could lead to cellular dysregulation, e.g, generation of an autocrine signaling loop.**
- **One subject (Subject #4) of ten successfully treated in Paris with retroviral gene therapy for X-SCID has developed a malignant proliferation of T lymphoid cells, in which the retroviral vector appears to have transactivated the LMO2 proto-oncogene.**

## Nosology of the SAE

- Immunophenotype dissimilar to typical pediatric T-ALL, which are not usually TCR- $\gamma/\delta$  leukemias.
- Gene activation by retroviral insertion is not a common etiology of human leukemias.
- Therefore, phrases other than “leukemia” have been used to describe SAE, e.g., “leukemoid”, “leukemia-like”, “lymphoproliferation”.

# Characteristics of Leukemia

- ✓ • Clonal proliferation of lymphohematopoietic cells
- ✓ • Genetic alteration(s)
- ✓ • Malignant
  - ✓ — Population growth
  - ✓ — Impairment of normal hematopoiesis
  - ✓ — Infiltration of peripheral organs
- ✓ • Genomic instability

## **Goals of Monitoring Protocol**

- 1) provide adequate monitoring so that determination of the need for therapeutic intervention is made as expeditiously as possible;**
- 2) assure that subjects are not exposed to either physical or psychological harm from unnecessary interventions;**
- 3) characterize any leukemias occurring in gene therapy trials;**
- 4) determine whether the leukemia resulted from insertional mutagenesis;**
- 5) characterize the clonality of hematopoiesis after retroviral gene therapy of HSC.**

## **Features of Monitoring Protocol**

- 1) Routine prospective monitoring of clonality of integrants in lymphohematopoietic cells by LAM-PCR.**
- 2) Routine monitoring for clinical signs of abnormally growing populations of cells.**
- 3) Detailed characterization of peripheral and/or marrow populations if indicated by increasing predominance of retroviral integrant; evidence of abnormal hematopoiesis (clinical or laboratory).**
- 4) Decision to intervene based on clinical and laboratory evidence of leukemia.**
- 5) Lifetime monitoring.**

## **Monitoring of Clonality**

- 1) LAM-PCR Q 6months for vector integrants.**
- 2) Increasing oligoclonality or monoclonality: 2-fold increase over two successive timepoints to >20% of total transduced cells.**
- 3) If increasing oligoclonality or monoclonality: initiate clinical evaluation, including blood counts, chemistries, evaluation for lymphoproliferation.**
- 4) If no evidence of leukemia, then repeat LAM-PCR Q 3months for two years until evidence of stabilization or leukemia.**
- 5) Lifetime routine monitoring.**

## **Techniques for Characterization of Increasingly Prevalent or Abnormal Clones**

- 1) Sequencing of LAM-PCR product to determine site of integration.**
- 2) If abnormal hematopoietic population, then analyze immunophenotype to assign lineage.**
- 3) If lymphoid cells, then appropriate immunoreceptor gene analyses to determine and characterize clonality, e.g., BCR or TCR.**
- 4) Cytogenetic analyses.**
- 5) Characterization of peripheral invasion and interference with hematopoiesis, i.e., marrow sample as indicated.**

# Diagnostic Questions re Increasingly Prevalent or Abnormal Clones

Is clone malignant?

## Clinical behavior

Population growth

Impairment of normal hematopoiesis

Infiltration of peripheral organs

## Molecular and cellular abnormalities

Cytogenetic abnormality

Inference from integration site

Abnormal differentiation

# Diagnostic Decisions re Increasingly Prevalent or Abnormal Clones

Is clone differentiating normally?

## Clinical behavior

Population growth stable

Normal hematopoiesis

No evidence of block in differentiation

## Molecular and cellular abnormalities

Integrand present in cells of multiple lineages

Diversity of TCR, BCR rearrangements



# Insertional Oncogenesis and Clonality of Hematopoiesis after Gene Therapy - IC Issues

## Investigators

Fabio Candotti M.D., Ph.D.	NHGRI
Donald Kohn, M.D.	CHLA
Greg Podsakoff, M.D.	CHLA
Cristof von Kalle, M.D.	U of Cincinnati
Kenneth Weinberg, M.D.	CHLA
Linda Forsyth, M.D.	FDA
Carolyn Wilson, Ph.D.	FDA
IRB members and staff	CHLA

## Nosology of the SAE

- Immunophenotype dissimilar to typical pediatric T-ALL, which are not usually TCR- $\gamma/\delta$  leukemias.
- Gene activation by retroviral insertion is not a common etiology of human leukemias.
- Therefore, phrases other than “leukemia” have been used to describe SAE, e.g., “leukemoid”, “leukemia-like”, “lymphoproliferation”.

# Characteristics of Leukemia

- ✓ • Clonal proliferation of lymphohematopoietic cells
- ✓ • Genetic alteration(s)
- ✓ • Malignant
  - ✓ — Population growth
  - ✓ — Impairment of normal hematopoiesis
  - ✓ — Infiltration of peripheral organs
- ✓ • Genomic instability

## Considerations in IC Discussion of Risk of Leukemia

- Discussion of incidence of leukemia would be desirable as means of presenting likely risk.
- However,
  1. The actual incidence of leukemia in gene therapy studies is unknown.
  2. It is not known what the relevant denominator for such a statement would be: the number of subjects in HSC gene therapy trials, the number in SCID trials, the number in X-SCID trials?
- Quantitative statement re risk would not be based on fact and could be coercive or misleading.